ISMP's List of *High-Alert* Medications

High-alert medications are drugs that bear a heightened risk of causing significant patient harm when they are used in error. Although mistakes may or may not be more common with these drugs, the consequences of an error are clearly more devastating to patients. We hope you will use this list to determine which medications require special safeguards to reduce the risk of errors. This may include strategies such as standardizing the ordering, storage, preparation, and administration of these products; improving access to information about these drugs; limiting access to high-alert medications; using auxiliary labels and automated alerts; and employing redundancies such as automated or independent double-checks when necessary. (Note: manual independent double-checks are not always the optimal error-reduction strategy and may not be practical for all of the medications on the list).

Classes/Categories of Medications
adrenergic agonists, IV (e.g., **EPINEPH**rine, phenylephrine, norepinephrine)
adrenergic antagonists, IV (e.g., propranolol, metoprolol, labetalol)
anesthetic agents, general, inhaled and IV (e.g., propofol, ketamine)
antiarrhythmics, IV (e.g., lidocaine, amiodarone)
antithrombotic agents, including: ■ anticoagulants (e.g., warfarin, low-molecular-weight heparin, IV unfractionated heparin) ■ Factor Xa inhibitors (e.g., fondaparinux) ■ direct thrombin inhibitors (e.g., argatroban, bivalirudin, dabigatran etexilate, lepirudin) ■ thrombolytics (e.g., alteplase, reteplase, tenecteplase) ■ glycoprotein IIb/IIIa inhibitors (e.g., eptifibatide)
cardioplegic solutions
chemotherapeutic agents, parenteral and oral
dextrose, hypertonic, 20% or greater
dialysis solutions, peritoneal and hemodialysis
epidural or intrathecal medications
hypoglycemics, oral
inotropic medications, IV (e.g., digoxin, milrinone)
insulin, subcutaneous and IV
liposomal forms of drugs (e.g., liposomal amphotericin B) and conventional counterparts (e.g., amphotericin B desoxycholate)
moderate sedation agents, IV (e.g., dexmedetomidine, midazolam)
moderate sedation agents, oral, for children (e.g., chloral hydrate)
narcotics/opioids ■ IV ■ transdermal ■ oral (including liquid concentrates, immediate and sustained-release formulations)
neuromuscular blocking agents (e.g., succinylcholine, rocuronium, vecuronium)
parenteral nutrition preparations
radiocontrast agents, IV
sterile water for injection, inhalation, and irrigation (excluding pour bottles) in containers of 100 mL or more
sodium chloride for injection, hypertonic, greater than 0.9% concentration

Specific Medications
epoprostenol (Flolan), IV
magnesium sulfate injection
methotrexate, oral, non-oncologic use
opium tincture
oxytocin, IV
nitroprusside sodium for injection
potassium chloride for injection concentrate
potassium phosphates injection
promethazine, IV
vasopressin, IV or intraosseous

Background

Based on error reports submitted to the ISMP National Medication Errors Reporting Program, reports of harmful errors in the literature, and input from practitioners and safety experts, ISMP created and periodically updates a list of potential high-alert medications. During October 2011-February 2012, 772 practitioners responded to an ISMP survey designed to identify which medications were most frequently considered high-alert drugs by individuals and organizations. Further, to assure relevance and completeness, the clinical staff at ISMP, members of our advisory board, and safety experts throughout the US were asked to review the potential list. This list of drugs and drug categories reflects the collective thinking of all who provided input.

© ISMP 2012. Permission is granted to reproduce material with proper attribution for internal use within healthcare organizations. Other reproduction is prohibited without written permission from ISMP. Report actual and potential medication errors to the ISMP National Medication Errors Reporting Program (ISMP MERP) via the website (www.ismp.org) or by calling 1-800-FAIL-SAF(E).